

HOMEWORK SET #3

Due: Tuesday, November 9

1. Consider the following scenarios of one-locus selection with two alleles, A and a and selection coefficient $s = 0.1$ against a . The genetic basis of fitness is given for each scenario. In each case, compute Δp , the between-generation change in the frequency of A for each of two initial frequencies: $p = 0.05$ and $p = 0.9$.
 - (a) haploid ($w_A = 1, w_a = 1-s$)
 - (b) diploid; A dominant advantageous
 - (c) diploid; A recessive advantageous
 - (d) diploid; A advantageous and partially dominant with $h = 0.4$
 - (e) Using your results, briefly discuss how the individual and population genetic architectures of fitness affect rates of adaptive evolution.

2. Consider a biallelic locus with alleles A and a and frequencies $p = 0.7$ and $q = 0.3$ respectively. Assume that prior to selection (i.e., in the zygote stage) the population is in Hardy-Weinberg proportions.
 - First consider “additive” fitnesses, i.e., $w_{AA} : w_{Aa} : w_{aa} = 1 : 1 - (s/2) : 1 - s$.
 - (a) Assuming $s = 0.2$, compute P_{AA}^* , P_{Aa}^* , and P_{aa}^* , the genotype frequencies after selection but before reproduction. Carry out your calculations to 4 or 5 decimal places.
 - (b) Show that P_{AA}^* , P_{Aa}^* , P_{aa}^* are not in Hardy-Weinberg proportions.
 - Now consider “multiplicative” fitnesses, i.e., $w_{AA} : w_{Aa} : w_{aa} = 1 : 1 - s : (1 - s)^2$.
 - (c) Redo part (a) using multiplicative fitnesses.
 - (d) Show that the post-selection genotype frequencies (P_{AA}^* , P_{Aa}^* , P_{aa}^*) are *exactly* in Hardy-Weinberg proportions.
 - (e) Explain why failure to reject the Hardy-Weinberg hypothesis statistically need not imply that a population satisfies the Hardy-Weinberg conditions.
 - (f) Show that with multiplicative fitnesses the evolutionary rate of p is, for any s ,

$$\Delta p = pq \frac{s}{1 - sq}$$
 (Note: this is identical to Δp for an advantageous allele in an *asexual* or *haploid* population with relative fitnesses $w_A = 1; w_a = 1 - s$.)

3.
 - (a) How large does the selection coefficient s need to be in order to change a dominant advantageous allele from an allele frequency of 0.5 to 0.51 in one generation?
 - (b) Do the same calculation for a recessive advantageous allele.
 - (c) Do the same calculation for a haploid advantageous allele.
 - (d) What does this imply about the efficiency of selection in haploids vs. diploids?
 - (Optional) Although this problem suggests that selection in haploids is more efficient than in diploids (Hint! Hint!), we know from problem 2 that selection in diploids can be equally efficient as haploids with multiplicative fitnesses (Triple Hint!). Are there any selection patterns in diploids that are more efficient than haploids for a given selection coefficient?

4. Hedrick, p. 362 #1

5. Hedrick, p. 362, #4